

Pyricularia grisea (oryzae) Modelo basado en Yoshino 1974, 1979 y Hashimoto et al 1984

Sensores:

T^a del aire

Humectación



Lluvia

Comienza si:

$$18 \leq (\sum_{1..120} T_{\text{hora}}) / 120 \leq 25$$

$$\text{Lluvia}_{\text{hora}} < 4\text{mm}$$

Humectación = Si



Infection finished following the Graph above (Yoshino 1979). Infection Severity is estimated by the average temperature during the leaf wetness period leading to infection following the second graph above.

Model Output:

Progress for Infection on Level1 and Level2 (0 - 100)

Infection severity (0 - 3)

Sheath Blight of rice (*Rhizoctonia solani*)

Simple risk model

Check last 120 hours:

If consecutive leaf wetness accumulate temperature depending values for every minute:

(12°C - 15°C = 1, 16°C - 17°C = 2, >18°C = 4)

If leaf wetness ends evaluate accumulated values:

Value > 4096 = RiskValue + 64 Points, Value - 4096

Value > 2048 = RiskValue + 16 Points, Value - 2048

Value > 1024 = RiskValue + 4 Points, Value - 1024

If global radiation is consecutive higher than 800 W/m² accumulate time in minutes and if radiation becomes lower evaluate values:

Value > 1024 = RiskValue - 32 Points, Value - 1024

Value > 512 = RiskValue - 8 Points, Value - 512

Value > 256 = RiskValue - 2 Points, Value - 256

Rice Blast Infection (*Pyricularia grisea (oryzae)*) Infection Model based on Yoshino 1974, 1979 and Hashimoto et al 1984 and Sheath Blight of rice (*Rhizoctonia solani*) simple risk model

Practical Use:

The rice blast model points out infection date and infection severity. This information is valid to improve the applied spray program. In a preventative spray program plant growth is reducing the protected plant surface within days. An infection which takes place immediately after a preventative spray will be covered perfectly. If an infection will be 6 or more days after the last spray disease control can be sustainably reduced by the high volume of uncovered leaves. In this case a curative spray will be indicated to ensure the success of disease control.

The sheath blight model points out periods with a high risk for this disease. No sprays will have to be applied in periods where the risk is low. In periods with moderate risk spray interval can be prolonged and in periods with high risk spray interval may have to be reduced or more effective compounds will have to be used.